

## Chapter 27 - Bacteria and Archaea

### Overview: Masters of Adaptation

- California's Owens Lake has a salt concentration of 32%, nine times saltier than seawater.
- Despite its harsh conditions, the lake's distinctive pink color is caused by **bacteriorhodopsin**, a photosynthetic pigment produced by trillions of *Halobacteria*, a single-celled archaean.
- This archaean is among the most salt-tolerant organisms on Earth. It pumps  $K^+$  into its cell until the ionic concentration within the cell matches the external salt concentration.
- Many other prokaryotes are adapted to extremely harsh conditions.
  - *Deinococcus radiodurans* can survive a radiation dose of 3,000,000 rads, while 1000 rads is fatal to a human.
  - *Picrophilus oshimae* can grow at a pH of 0.03, acidic enough to dissolve metal.
  - Some prokaryotes live in rocks 3.2 kilometers below the Earth's surface.
- Prokaryotes are adapted to a broad range of habitats, including the land and waters in which other species are found.
- Today, prokaryotes still dominate the biosphere.
  - Their collective biomass outweighs that of all eukaryotes combined by at least tenfold.
  - More prokaryotes inhabit a handful of fertile soil or the mouth or skin of a human than the total number of people who have ever lived.

### Concept 27.1 Structural and functional adaptations contribute to prokaryotic success.

*Prokaryotes are small.*

- Prokaryotes were the first organisms to live on Earth.
- Most prokaryotes are unicellular.
- Some species may aggregate transiently or permanently in colonies.
- Most prokaryotes have diameters in the range of 0.5–5  $\mu$ m, compared to 10–100  $\mu$ m for most eukaryotic cells.
  - The largest prokaryote discovered so far has a diameter of 750  $\mu$ m, just visible to the unaided eye.
- The most common shapes among prokaryotes are spheres (cocci), rods (bacilli), and spirals.

*Nearly all prokaryotes have a cell wall external to the plasma membrane.*

- In nearly all prokaryotes, a cell wall maintains the shape of the cell, affords physical protection, and prevents the cell from bursting in a hypotonic environment.
- In a hypertonic environment, most prokaryotes lose water and plasmolyze, like other walled cells.
  - Severe water loss inhibits the reproduction of prokaryotes, which explains why salt can be used to preserve foods.
- Most bacterial cell walls contain **peptidoglycan**, a polymer of modified sugars cross-linked by short polypeptides.
  - This molecular fabric encloses the entire bacterium and anchors other molecules that extend from its surface.
  - The cell walls of archaea contain polysaccharides and proteins, but lack peptidoglycan.
- The **Gram stain** is a valuable tool for identifying specific bacteria based on differences in their cell walls.
  - **Gram-positive** bacteria have simple cell walls with large amounts of peptidoglycans.
  - **Gram-negative** bacteria have more complex cell walls with less peptidoglycan.
- An outer membrane on the cell wall of gram-negative cells contains lipopolysaccharides, carbohydrates bonded to lipids.
- Among pathogenic bacteria, gram-negative species are generally more deadly than gram-positive species.
  - The lipopolysaccharides on the walls of gram-negative bacteria are often toxic, and the outer membrane protects the pathogens from the defenses of their hosts.
  - Gram-negative bacteria are commonly more resistant than gram-positive species to antibiotics because the outer membrane impedes entry of the drugs.
- Many antibiotics, including penicillin, inhibit the synthesis of cross-links in peptidoglycans, preventing the formation of a functional wall, especially in gram-positive species.
  - These drugs cripple many species of bacteria, without affecting human and other eukaryote cells that do not synthesize peptidoglycans.
- Many prokaryotes secrete another sticky protective layer of polysaccharide or protein, the **capsule**, outside the cell wall.
  - Capsules allow cells to adhere to their substrate or to other individuals in a colony.
  - Some capsules protect against dehydration, and some may increase resistance to host defenses.
- Another way for prokaryotes to adhere to one another or to the substratum is by surface appendages called **fimbriae**, also known as *attachment pili*.
  - Fimbriae are usually more numerous and shorter than **sex pili**.
  - Sex pili are specialized for holding two prokaryote cells together long enough to transfer DNA during conjugation.

*Many prokaryotes are motile.*

- About half of all prokaryotes are capable of directional movement.
  - Some species can move at speeds exceeding 50 m/sec, about 50 times their body length per second.
- The beating of flagella scattered over the entire surface or concentrated at one or both ends is the most common method of movement.
  - The flagella of prokaryotes differ in structure and function from those of eukaryotes.
- In a heterogeneous environment, many prokaryotes are capable of **taxis**, movement *toward* nutrients or oxygen (positive chemotaxis) or *away from* a toxic substance (negative chemotaxis).
  - Prokaryotes that exhibit chemotaxis respond to chemicals by changing their movement patterns.
  - Solitary *Escherichia coli* may exhibit positive chemotaxis toward other members of their species, enabling the formation of colonies.

*The cellular and genomic organization of prokaryotes is fundamentally different from that of eukaryotes.*

- The cells of prokaryotes are simpler than those of eukaryotes in both internal structure and genomic organization.
- Prokaryotic cells lack the complex compartmentalization found in eukaryotic cells.
- Instead, prokaryotes use specialized infolded regions of the plasma membrane to perform many metabolic functions, including cellular respiration and photosynthesis.
- Prokaryotes have smaller, simpler genomes than eukaryotes.
- In the majority of prokaryotes, the genome consists of a ring of DNA with few associated proteins.
- The prokaryotic chromosome is located in the **nucleoid region**.
- Prokaryotes may also have smaller rings of DNA called **plasmids**, which consist of only a few genes.
- Although the general processes for DNA replication and translation of mRNA into proteins are fundamentally alike in eukaryotes and prokaryotes, some of the details differ.
  - For example, prokaryotic ribosomes are slightly smaller than the eukaryotic version and differ in protein and RNA content.
- These differences are great enough that selective antibiotics, including tetracycline and erythromycin, bind to prokaryotic ribosomes to block protein synthesis in prokaryotes but not in eukaryotes.

*Populations of prokaryotes grow and adapt rapidly.*

- Prokaryotes have the potential to reproduce quickly in a favorable environment.
  - While most prokaryotes have generation times of 1–3 hours, some species can produce a new generation in 20 minutes under optimal conditions.
  - A single cell in favorable conditions produces a large colony of offspring very quickly.

- Prokaryotes reproduce asexually via **binary fission**, synthesizing DNA almost continuously.
- Prokaryotic reproduction is limited because cells eventually exhaust their nutrient supply, accumulate metabolic wastes, face competition from other microbes, or are consumed by other organisms.
- Some bacteria form resistant cells called **endospores** when an essential nutrient is lacking in the environment.
  - A cell replicates its chromosome and surrounds one chromosome with a durable wall to form the endospore. Water is removed from the endospore, halting metabolism.
  - The original cell then disintegrates to leave the endospore behind.
- An endospore is resistant to all sorts of trauma.
  - Most endospores can survive in boiling water.
  - Endospores may remain dormant but viable for centuries or longer.
  - When the environment becomes more hospitable, the endospore absorbs water and resumes growth.
  - Sterilization in an autoclave kills endospores by heating them to 121°C under high pressure.
- Mutation is the major source of genetic variation in prokaryotes.
- With generation times of minutes or hours, prokaryotic populations can adapt very rapidly to environmental changes as natural selection favors gene mutations that confer greater fitness.
- As a consequence, prokaryotes are important model organisms for scientists who study evolution in the laboratory.
- Prokaryotes are highly evolved. For more than 3.5 billion years, prokaryotic populations have responded successfully to many different types of environmental challenge.
- Prokaryotic populations harbor high levels of genetic diversity on which selection can act.

**Concept 27.2 Rapid reproduction, mutation, and genetic recombination promote genetic diversity in prokaryotes.**

- Prokaryotic populations contain considerable genetic variation.
  - For example, a ribosomal RNA gene differs more between two strains of *E. coli* than it does between a human and a platypus.
- Three factors give rise to high levels of genetic diversity in prokaryotes: rapid reproduction, mutation, and genetic recombination.
- When a prokaryote reproduces by binary fission, some of the offspring differ slightly in genetic makeup due to mutation.
- The probability of a spontaneous mutation in a given *E. coli* gene is only about  $1 \times 10^{-7}$  per cell division. However, among the  $2 \times 10^{10}$  new *E. coli* cells that arise each day in a single human colon, approximately 2,000 will have a mutation in that gene.
  - When all 4,300 *E. coli* genes are considered, 9 million mutant *E. coli* cells arise per day per

human host.

- The genetic diversity within a species like *E. coli* can lead to rapid evolution, as cells that are better equipped for the local environment survive and reproduce more successfully than others.
- Genetic recombination is another factor that generates diversity within bacterial populations.
  - Here, *recombination* is defined as the combining of DNA from two individuals into a single genome.
- Bacterial recombination occurs through three processes: transformation, transduction, and conjugation.
- **Transformation** is the alteration of a bacterial cell's genotype by the uptake of naked, foreign DNA from the surrounding environment.
  - For example, harmless *Streptococcus pneumoniae* bacteria can be transformed to pneumonia-causing cells.
  - This transformation occurs when a live nonpathogenic cell takes up a piece of DNA that happens to include the allele for pathogenicity from dead, broken-open pathogenic cells.
  - The foreign allele replaces the native allele in the bacterial chromosome by genetic recombination, with an exchange of homologous DNA segments.
  - The resulting cell is now recombinant, with DNA derived from two different cells.
- Years after transformation was discovered in laboratory cultures, most biologists believed that the process was too rare and haphazard to play an important role in natural bacterial populations.
- Researchers have since learned that many bacterial species have surface proteins that are specialized for the uptake of naked DNA.
- These proteins recognize and transport DNA from closely related bacterial species into the cell, which can then incorporate the foreign DNA into the genome.
- **Transduction** is a type of horizontal gene transfer that occurs when a phage carries bacterial genes from one host cell to another, as a result of aberrations in the phage reproductive cycle.
- Occasionally, a small piece of the host cell's degraded DNA, rather than the phage genome, is packaged within a phage capsid.
  - When this phage attaches to another bacterium, it injects this foreign DNA into its new host.
  - Some of this DNA can subsequently replace the homologous region of the second cell.
- Sometimes known as bacterial "sex," **conjugation** transfers genetic material between two bacterial cells that are temporarily joined.
- The transfer is one-way. One cell donates DNA, and its "mate" receives the genes.
- A hollow sex pilus from the donor initially joins the two cells and retracts to pull the two cells together.

- A temporary *mating bridge* forms between the cells.
- The ability to form a sex pilus and donate DNA during conjugation results from an **F factor** (F for *fertility*) as a section of the bacterial chromosome or as a plasmid.
  - The F factor consists of about 25 genes, most required for the production of sex pili.
- A cell containing the **F plasmid** is designated an  $F^+$  cell and functions as a DNA donor during conjugation.
  - A cell lacking the F factor, designated an  $F^-$  cell, functions as a DNA recipient.
  - Transfer of the  $F^+$  plasmid converts an  $F^-$  cell to an  $F^+$  cell.
- The F factor can become integrated into the bacterial chromosome.
- A cell with the F factor built into its chromosome is called an Hfr cell (Hfr for *high frequency of recombination*).
- Hfr cells function as donors during conjugation.
  - When chromosomal DNA from an Hfr cell enters an  $F^-$  cell, homologous regions of the Hfr and  $F^-$  chromosomes may align, allowing segments of their DNA to be exchanged.
  - The result is the production of a recombinant bacterium that has genes derived from two different cells.
- Although the processes of horizontal gene transfer have been studied only in bacteria, it is assumed that they are also important in archaea.
- In the 1950s, Japanese physicians began to notice that some bacterial strains had evolved antibiotic resistance.
  - A mutation may reduce the ability of the pathogen's cell-surface proteins to transport a particular antibiotic into the bacterial cell.
  - A mutation in a different gene may alter the intracellular target protein for an antibiotic molecule, reducing its effect.
  - Some bacteria have resistance genes coding for enzymes that specifically destroy certain antibiotics, like tetracycline or ampicillin.
- The genes conferring resistance are carried by plasmids, specifically the **R plasmid** (R for *resistance*).
- When a bacterial population is exposed to an antibiotic, individuals with the R plasmid survive and increase in the overall population.
- Because R plasmids also have genes that encode for sex pili, they can be transferred from one cell to another by conjugation.
- Some R plasmids carry as many as ten genes for resistance to ten different antibiotics.

**Concept 27.3 A great diversity of nutritional and metabolic adaptations have evolved in prokaryotes.**

- Organisms can be categorized by their nutrition, based on how they obtain energy and carbon to build the organic molecules that make up their cells.
- Nutritional diversity is greater among prokaryotes than among all eukaryotes.
- Every type of nutrition observed in eukaryotes is found in prokaryotes, along with some nutritional modes unique to prokaryotes.
  - Organisms that obtain energy from light are *phototrophs*.
  - Organisms that obtain energy from chemicals in their environment are *chemotrophs*.
  - Organisms that need only an inorganic compound such as CO<sub>2</sub> as a carbon source are *autotrophs*.
  - Organisms that require at least one organic nutrient—such as glucose—as a carbon source are *heterotrophs*.
- These categories of energy source and carbon source can be combined to group prokaryotes according to four major modes of nutrition.
- **Photoautotrophs** are photosynthetic organisms that harness light energy to drive the synthesis of organic compounds from CO<sub>2</sub> or other inorganic carbon compounds such as HCO<sub>3</sub><sup>-</sup>.
  - Among the photoautotrophic prokaryotes are the cyanobacteria.
  - Among the photosynthetic eukaryotes are plants and algae.
- **Chemoautotrophs** need only an inorganic molecule like CO<sub>2</sub> as a carbon source but obtain energy by oxidizing inorganic substances.
  - These inorganic substances include hydrogen sulfide (H<sub>2</sub>S), ammonia (NH<sub>3</sub>), and ferrous ions (Fe<sup>2+</sup>), among others.
  - This nutritional mode is unique to prokaryotes.
- **Photoheterotrophs** use light to generate ATP but obtain their carbon in organic form.
  - This mode of nutrition is restricted to a few marine and halophilic prokaryotes.
- **Chemoheterotrophs** must consume organic molecules for both energy and carbon.
  - This nutritional mode is found widely in prokaryotes, protists, fungi, animals, and even some parasitic plants.
- Prokaryotic metabolism also varies with respect to oxygen.
  - **Obligate aerobes** require O<sub>2</sub> for cellular respiration.
  - **Facultative anaerobes** use O<sub>2</sub> if it is present but can also grow by fermentation in an anaerobic environment.
  - **Obligate anaerobes** are poisoned by O<sub>2</sub> and use either fermentation or **anaerobic respiration**, in which inorganic molecules other than O<sub>2</sub> accept electrons from electron transport chains.

- Diverse prokaryotes can metabolize a wide variety of compounds that contain nitrogen, an essential component of proteins and nucleic acids in all organisms.
  - Eukaryotes are limited in the forms of nitrogen they can use.
- **Nitrogen-fixing** prokaryotes convert  $N_2$  to  $NH_3$ , making atmospheric nitrogen available to themselves (and eventually to other organisms) for incorporation into organic molecules.
- Nitrogen-fixing cyanobacteria are the most self-sufficient of all organisms.
  - These cyanobacteria require only light energy,  $CO_2$ ,  $N_2$ , water, and some minerals to grow.
- Prokaryotes were once thought of as single-celled individualists.
- Microbiologists now recognize that cooperation between prokaryotes allows them to use environmental resources they cannot exploit as individuals.
- Cooperation may involve specialization in the cells of a prokaryotic colony.
  - For example, the cyanobacterium *Anabaena* forms filamentous colonies with specialized cells to carry out nitrogen fixation.
  - Photosynthesis produces  $O_2$ , which inactivates the enzymes involved in nitrogen fixation.
  - Most cells in the filament are photosynthetic, while a few specialized cells called **heterocytes** carry out only nitrogen fixation.
  - A heterocyte is surrounded by a thickened cell wall that restricts the entry of oxygen produced by neighboring photosynthetic cells.
  - Heterocyte transport fixes nitrogen to neighboring cells in exchange for carbohydrates.
- In some prokaryotic species, metabolic cooperation occurs in surface-coating colonies known as **biofilms**.
  - Cells in a biofilm secrete signaling molecules to recruit nearby cells, causing the colony to grow.
  - Once the colony is sufficiently large, the cells begin producing proteins that adhere the cells to the substrate and to one another.
  - Channels in the biofilms allow nutrients to reach cells in the interior and allow wastes to be expelled.
- In some cases, different species of prokaryotes may cooperate.
  - For example, sulfate-consuming bacteria and methane-consuming archaea coexist in ball-shaped aggregates in the mud of the ocean floor.
  - The bacteria use the archaea's waste products, such as organic compounds and hydrogen.
  - In turn, the bacteria produce compounds that facilitate methane consumption by the archaea.
  - Each year, these archaea consume an estimated 300 billion kg of methane, a major greenhouse gas.

#### **Concept 27.4 Molecular systematics is illuminating prokaryotic phylogeny.**

- Until the late 20th century, systematists based prokaryotic taxonomy on criteria such as shape, motility, nutritional mode, and Gram staining.
- Although these criteria may be valuable in culturing and identifying pathogenic bacteria, they may not reflect evolutionary relationships.

- Applying molecular systematics to the investigation of prokaryotic phylogeny has been very fruitful.
- Microbiologists began comparing sequences of prokaryotic genes in the 1970s.
  - Carl Woese and his colleagues used small-subunit ribosomal RNA (SSU-rRNA) as a marker for evolutionary relationships.
  - They concluded that many prokaryotes once classified as bacteria are actually more closely related to eukaryotes and that they belong in a domain of their own—Archaea.
- Microbiologists have since analyzed larger amounts of genetic data, including whole genomes of some species.
  - They found that a few traditional taxonomic groups, such as cyanobacteria, are monophyletic.
  - Other groups, such as gram-negative bacteria, are scattered throughout several lineages.
- One important lesson that has already emerged from studies of prokaryotic phylogeny is that the genetic diversity of prokaryotes is immense.
- When researchers began to sequence the genes of prokaryotes, they could investigate only those species that can be cultured in the laboratory, a tiny minority of all prokaryotes.
- Norman Pace of the University of Colorado pioneered the use of the polymerase chain reaction (PCR) to analyze the genes of prokaryotes that are collected directly from their environment, such as water or soil samples.
  - Such “genetic prospecting” is now widely used.
- Every year, new prokaryotes are identified that add major new branches to the tree of life.
  - Some researchers suggest that certain branches represent new kingdoms.
- Although only 6,300 prokaryotes have been fully characterized, a single handful of soil could contain 10,000 prokaryotic species, according to some estimates.
- Another important lesson is the significance of horizontal gene transfer in the evolution of prokaryotes.
- Over hundreds of millions of years, prokaryotes have acquired genes from distantly related species, and they continue to do so today.
- As a result, significant portions of the genomes of many prokaryotes are actually mosaics of genes imported from other species.
- Horizontal gene transfer obscures the location of the root of the tree of life.
- Still, it is clear that very early in the history of life on Earth, prokaryotes diverged into two main lineages: bacteria and archaea.

*Researchers are identifying a great diversity of archaea in extreme environments and in the oceans.*

- Archaea share certain traits with bacteria and other traits with eukaryotes.
- Archaea also have many unique characteristics, as expected for a taxon that has followed a separate evolutionary path for so long.

- Much of the research on archaea has focused not on phylogeny but on their ecology—their ability to live where no other life can.
- The first prokaryotes to be classified in domain Archaea are species that can live in environments so extreme that few other organisms can survive there.
- Such organisms are known as **extremophiles** or “lovers” of extreme environments.
- Extremophiles include extreme thermophiles and extreme halophiles.
- **Extreme halophiles** live in such salty places as the Great Salt Lake, the Dead Sea, and Owens Lake.
  - Some species merely tolerate elevated salinity; others require an extremely salty environment to grow.
  - For example, the proteins and cell walls of *Halobacteria* have features that function only at high salinity. These organisms cannot live below 9% salinity.
- **Extreme thermophiles** thrive in hot environments.
  - The optimum temperatures for most thermophiles are 60–80°C.
  - *Sulfolobus* oxidizes sulfur in 90°C sulfur springs in Yellowstone National Park.
  - Another sulfur-metabolizing thermophile can double its cell numbers in 24 hours at temperatures as high as 121°C in water near deep-sea hydrothermal vents.
  - *Pyrococcus furiosus* is an extreme thermophile that is used in biotechnology as the source of DNA polymerase for the polymerase chain reaction (PCR).
- Other archaea do not live in extreme environments.
- **Methanogens** obtain energy by using CO<sub>2</sub> to oxidize H<sub>2</sub>, producing methane as a waste product.
  - Methanogens are among the strictest anaerobes and are poisoned by O<sub>2</sub>.
  - Although some methanogens live in extreme environments, such as buried under kilometers of ice in Greenland, other species live in swamps and marshes where other microbes have consumed all the oxygen.
    - ♣ “Marsh gas” is actually methane produced by archaea.
  - Other methanogens live in the anaerobic guts of animals, playing an important role in their nutrition.
  - Methanogens are important decomposers in sewage treatment facilities.
- Many extreme halophiles and all known methanogens, plus a few extreme thermophiles, are members of a clade called Euryarchaeota.
- Most thermophilic species belong to a second clade, Crenarchaeota.
- Genetic prospecting has revealed that both Euryarchaeota and Crenarchaeota include many species of archaea that are not extremophiles.
  - These species exist in habitats ranging from farm soils to lake sediments to the surface waters of the ocean.
- New findings continue to update our understanding of archaean phylogeny.
- A new clade, Korarchaeota, has been identified that appears to be the oldest lineage in the domain Archaea.

- In 2002, researchers exploring hydrothermal vents off the coast of Iceland discovered archaean cells only 0.4 μm in diameter attached to a much larger crenarchaeote.
- The genome of this tiny archaean is one of the smallest known of any organisms, containing only 500,000 base pairs.
- This prokaryote belongs to a fourth archaean clade called Nanoarchaeota.
  - Three new nanoarchaeote species have since been found, one from Yellowstone's hot springs, one from hot springs in Siberia, and one from a hydrothermal vent in the Pacific.

*Bacteria include the vast majority of familiar prokaryotes.*

- Bacteria range from the pathogenic species that cause strep throat to the beneficial species that make Swiss cheese.
- Every major mode of metabolism and nutrition is represented among bacteria, which have a great impact on Earth and its life.

### **Concept 27.5 Prokaryotes play crucial roles in the biosphere.**

- If humans were to disappear from the planet tomorrow, life on Earth would go on for most other species.
- Prokaryotes are so important to the biosphere, however, that if they were to disappear, the prospects for any other life surviving would be dim.
- The atoms that make up the organic molecules in all living things were at one time part of inorganic compounds in the soil, air, and water.
- Life depends on the recycling of chemical elements between the biological and chemical components of ecosystems. Prokaryotes play an important role in this process.
  - Chemoheterotrophic prokaryotes function as **decomposers**, breaking down corpses, dead vegetation, and waste products and unlocking supplies of carbon, nitrogen, and other elements essential for life.
- Prokaryotes convert inorganic compounds into forms that can be taken up by other organisms.
  - Autotrophic prokaryotes use CO<sub>2</sub> to make organic compounds, which are then passed up through food chains.
  - Cyanobacteria produce atmospheric O<sub>2</sub>, and a number of prokaryotes fix atmospheric nitrogen (N<sub>2</sub>) into a form that other organisms can use to make proteins and nucleic acids.
- Prokaryotes may act to *increase* or *decrease* the availability of key plant nutrients. As a result, they have large but complex effects on soil nutrient concentrations.
- In marine environments, prokaryotes from the clade dominate the oceans with an estimated 10<sup>28</sup> cells.
  - These abundant organisms perform nitrification and may have a large impact on the global nitrogen cycle.

- Prokaryotes often interact with other species of prokaryotes or eukaryotes with complementary metabolisms.
- An ecological relationship between organisms that are in direct contact is called **symbiosis**.
- If one of the symbiotic organisms is larger than the other, it is called the **host**, and the smaller is known as the **symbiont**.
- In **commensalism**, one symbiotic organism benefits while the other is neither harmed nor helped by the relationship.
  - For example, more than 150 bacterial species live on the surface of your body, at densities of up to 10 million cells per  $\text{cm}^2$ . Most are commensalists.
- In **parasitism**, one symbiotic organism, the parasite, benefits at the expense of the host.
  - The **parasite** eats the cell contents, tissues, or body fluids of the host. Unlike predators, parasites do not kill the host, at least not immediately.
  - Parasites that cause disease are called **pathogens**.
- In **mutualism**, both symbiotic organisms benefit.
- Human intestines are home to an estimated 500 to 1,000 species of bacteria, which outnumber all human cells in the body tenfold.
  - Many of these species are mutualists, digesting food that our own intestines cannot.
- In 2003, scientists at Washington University in St. Louis published the first complete genome for a gut mutualist, *Bacteroides thetaiotaomicron*.
  - The genome includes a large array of genes involved in synthesizing carbohydrates, vitamins, and other nutrients needed by humans.
  - Signals from the bacterium activate human genes that build the network of intestinal blood vessels necessary to absorb food.
  - Other signals induce human cells to produce antimicrobial compounds to which *B. thetaiotaomicron* is not susceptible, protecting the bacterium from its competitors.

### **Concept 27.6 Prokaryotes have both harmful and beneficial impacts on humans.**

- Pathogenic prokaryotes represent only a small fraction of prokaryotic species.
- Prokaryotes cause about half of human diseases.
- Approximately 2 million people a year die of the lung disease tuberculosis, caused by the bacillus *Mycobacterium tuberculosis*.
- Another 2 million die from diarrhea caused by other prokaryotes.
- Lyme disease, caused by a bacterium carried by ticks that live on deer and field mice, is the most widespread pest-carried disease in the United States.
  - If untreated, Lyme disease can lead to debilitating arthritis, heart disease, and nervous disorders.

- Pathogens cause illness by producing poisons called exotoxins and endotoxins.
- **Exotoxins** are proteins secreted by prokaryotes.
- Exotoxins can produce disease symptoms even if the prokaryote is not present.
  - An exotoxin produced by *Vibrio cholerae* causes cholera, a serious disease characterized by severe diarrhea.
  - *V. cholerae* stimulates intestinal cells to release chloride ions ( $\text{Cl}^-$ ) into the gut; water follows by osmosis.
  - *Clostridium botulinum*, which grows anaerobically in improperly canned foods, produces an exotoxin that causes botulism.
- **Endotoxins** are lipopolysaccharide components of the outer membrane of some gram-negative bacteria.
- In contrast to exotoxins, endotoxins are released only when the bacteria die and their cell walls break down.
  - The endotoxin-producing bacteria in the genus *Salmonella* are not normally present in healthy animals.
  - *Salmonella typhi* causes typhoid fever.
  - Other *Salmonella* species, including some that are common in poultry, cause food poisoning.
- Since the discovery that “germs” cause disease, improved sanitation and improved treatments have reduced mortality and extended life expectancy in developed countries.
- Antibiotics have greatly reduced the threat of pathogenic prokaryotes and have saved a great many lives.
- However, resistance to antibiotics is currently evolving in many strains of prokaryotes.
- The rapid reproduction of prokaryotes enables genes conferring resistance to multiply quickly through prokaryotic populations as a result of natural selection.
- These genes can spread to other species by horizontal gene transfer.
- Horizontal gene transfer can also spread genes associated with virulence, turning harmless prokaryotes into fatal pathogens.
- *E. coli* is ordinarily a harmless symbiont in the human intestines, but pathogenic strains causing bloody diarrhea have arisen.
  - One of the most dangerous strains is called O157:H7.
  - Today, it is a global threat, with 75,000 cases annually in the United States alone.
- In 2001, an international team of scientists sequenced the genome of O157:H7 and compared it with the genome of a harmless strain of *E. coli*.
  - Of the 5,416 genes in O157:H7, 1,387 have no counterpart in the harmless strain.
  - These 1,387 genes must have been incorporated into the genome of O157:H7 through horizontal gene transfer, most likely through the action of bacteriophages.
  - Many of the imported genes are associated with the pathogen’s invasion of its host.
  - For example, some genes code for adhesive fimbriae that enable O157:H7 to attach itself to the intestinal wall and extract nutrients.
- Pathogenic prokaryotes pose a potential threat as weapons of bioterrorism.

- In October 2001, endospores of *Bacillus anthracis*, the bacterium that causes anthrax, were mailed to members of the news media and the U.S. Senate.
- Other prokaryotes that could be used as weapons include *C. botulinum* and *Yersinia pestis*, which causes plague.
- The threat of bioterrorism has stimulated intense research on pathogenic prokaryotes.
- Humans have learned to exploit the diverse metabolic capabilities of prokaryotes for scientific research and for practical purposes.
  - Humans have long used bacteria to make cheese and yogurt.
  - Our greater understanding of prokaryotes has led to many new biotechnology applications.
- Increasingly, prokaryotes are used to solve environmental problems.
- The use of organisms to remove pollutants from air, water, and soil is called **bioremediation**.
  - The most familiar example of bioremediation is the use of prokaryote decomposers to treat human sewage.
  - Anaerobic bacteria decompose the organic matter into material that can be used as landfill or fertilizer.
  - Other bioremediation applications include precipitating radioactive material from groundwater and cleaning up oil spills.
- Bacteria can be used to make durable, biodegradable natural plastics.
- Through genetic engineering, humans can now modify prokaryotes to produce vitamins, antibiotics, hormones, and many other products.
- Bioengineered prokaryotes can produce ethanol from agricultural waste, switchgrass, and fast-growing woody plants.
- Craig Venter of the Human Genome Project has announced that he and his colleagues are attempting to build “synthetic chromosomes” for prokaryotes, producing new species from scratch.
  - Venter hopes to “design” prokaryotes that can perform specific tasks, such as producing large amounts of hydrogen to reduce dependence on fossil fuels.
  - Other scientists question the wisdom of such an approach.
- The usefulness of prokaryotes derives from their diversity in metabolism and nutrition.